#### **Outpatients**


An outpatient's appointment will be arranged as requested by your consultant. It is important that your child attends his/her outpatient appointment, so that we can monitor his/her progress.

Outpatients Appointment
Date:
Time:



<u>Remember</u>: Ask your nurse if you are unsure about anything about your child's care.

# **Additional instructions**



Developed by Claire O'Byrne & staff of the Surgical Day Unit

Date of issue: April 2021 / Review Date: 2024

Disclaimer & Copyright ©2021, Children's Health Ireland at Crumlin, Dublin 12. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means without the prior written permission of the copyright holder. Every effort has been made to ensure that the information provided is accurate and in accord with standards accepted at the time of printing.



Children's Health Ireland at Crumlin

Phone: 01 409 6100

### What is a Knee Arthroscopy?

Knee Arthroscopy is a minor surgical procedure during which an arthroscope - a narrow tube with a light, is inserted into the knee joint. Your surgeon will carry out this procedure in the Operating Theatre and your child will have a general anaesthetic. It is usually performed as a day case. An Arthroscopy can assist in detecting an injury, damage or defect within the knee joint.

# After your child's Arthroscopy Pain Relief

- Local anaesthetic is usually given into the knee joint during the procedure to provide pain relief.
- Other pain relieving medications will be given to your child during surgery.
- Your nurse will give you a pain relief leaflet explaining the times your child is next due pain medication.
- Pain relief should be given at regular intervals until your child is comfortable and pain free.
- Distracting your child with their favourite play may also help take their mind off any pain or discomfort.



#### Wound Care

- On return to the ward, your child will have a pressure bandage on their knee. This bandage should remain in place for 48 hours, unless it becomes too tight or uncomfortable.
- Your child may have a brace on the knee that will be applied in theatre. It is called a Donjoy brace. If your child has one of these applied the Physiotherapist will discuss the management of this at home before you leave.
- If your child needs crutches, the physiotherapist will see your child prior to discharge.
- The small dressing beneath the pressure bandage should remain in place for 7-10 days.
- After 7-10 days, you may remove this small dressing.
- Steristrips (paper stitches) are usually present under the small dressing. These will fall off themselves. If they are still in place after 14 days, you may gently remove them.
- Keep all dressings dry until they are removed.





# Signs of infection

Please observe for signs of infection and seek medical attention if you are concerned.

These signs include:

- Excessive pain in the wound area
- Swelling in and around the wound site
- Redness around the wound site
- High temperature
- Yellow discharge/ oozing onto dressing

